

## **Pandas, Plants, and People**

Author(s): Jianguo Liu and Andrés Viña

Source: *Annals of the Missouri Botanical Garden*, 100(1-2):108-125. 2014.

Published By: Missouri Botanical Garden

URL: <http://www.bioone.org/doi/full/10.3417/2013040>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Use of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

# PANDAS, PLANTS, AND PEOPLE<sup>1,2</sup> *Jianguo Liu<sup>3</sup> and Andrés Viña<sup>3</sup>*

---

## ABSTRACT

Plants are essential for the survival and sustainability of both humans and wildlife species around the world. However, human activities have directly and indirectly affected almost all plants, which in turn have produced cascading effects on humans and wildlife through disruption of crucial ecosystem services and wildlife habitat. Understanding such complex interactions is crucial for developing better policies that reconcile the needs of an ever-growing human population with biodiversity conservation. Using the coupled human and natural systems (CHANS) framework, this article synthesizes research on the complex interactions of plant species, giant pandas, and people. The CHANS framework is particularly useful for uncovering key patterns and processes behind plant-animal interactions modified by human activities. Our synthesis shows that many human factors, including socioeconomic and demographic, together with other factors (e.g., projected global climate change), exhibit reciprocal interactions with pandas and the plant species that comprise their habitat. Although substantial efforts have been made to preserve plants and wildlife, much work still remains to be done, including the expansion and more effective management of protected areas, use of native plant species in reforestation/afforestation programs, and active participation of local residents in conservation actions.

*Key words:* Bamboo, China, coupled human and natural systems (CHANS), ecosystem services, endangered species, giant pandas, protected areas, wildlife.

---

Plants provide essential sources of food, fiber, medicine, and other important goods and services for humans (Díaz et al., 2006). They also constitute crucial components of wildlife habitats (Tuanmu et al., 2011), offering not only food to herbivorous wildlife species but also shelter to herbivorous and carnivorous wildlife species across many different ecosystems (Taylor et al., 2004; Nilsson & Wardle, 2005; Gilliam, 2007). However, human activities have converted much of the natural landscapes to human-dominated landscapes and have led to biodiversity loss worldwide (Vitousek et al., 1997; Sala et al., 2000; Pereira et al., 2004; Waltert et al., 2004; Lepczyk et al., 2008). This loss is usually expressed as species extinction at local, regional, and global scales and homogenization of regional and continental biotas. The rate of species extinction due to human activities is between 1000 and 10,000 times faster than background rates, i.e., those without human causes (Hilton-Taylor, 2000; Pimm & Raven, 2000; Ceballos et al., 2010). As a result, according to criteria of the International Union for Conservation of Nature (IUCN, 2001), estimates of the percentages of plant species worldwide that may be threatened with

extinction range from ca. 22% to as many as 50% (Pitman & Jørgensen, 2002; Baillie et al., 2004). The loss of plant biodiversity poses enormous direct and indirect consequences for humans as well as wildlife species and, ultimately, the structure and function of ecosystems (Hooper et al., 2005). Therefore, the loss of plant biodiversity is of great concern not only for ethical and aesthetic reasons but also for its cascading effects on entire ecosystems and the goods and services they provide to humanity.

The loss of biodiversity has inspired many conservation actions, such as the establishment of protected areas and implementation of conservation policies. Protected areas are supposed to be areas where human activities are limited or controlled (DeFries et al., 2007). By 2011, there were over 130,700 protected areas, covering more than 24 million km<sup>2</sup> of surface area on the earth (IUCN, UNEP-WCMC, 2012). Although their creation constitutes a cornerstone of biodiversity conservation (Margules & Pressey, 2000), biodiversity inside these areas is not necessarily better protected than outside them (Liu et al., 2001; Caro, 2002; Parks & Harcourt, 2002; Meir et al., 2004; Viña et al., 2007), because

---

<sup>1</sup> We thank numerous students and collaborators (especially Zhiyun Ouyang and Hemin Zhang) as well as the Wolong Administrative Bureau and Chinese Academy of Sciences for logistic assistance in field data collection and interviewees at Wolong Nature Reserve for their participation in this study. We are very grateful to Victoria C. Hollowell and Peter Raven for their invitation to write this paper, and to Peter Hoch, Victoria C. Hollowell, Maryjane Hotaling, and Jon Ricketson, for their very valuable insights and suggestions that improved the manuscript. We gratefully acknowledge financial support from agencies such as the National Science Foundation, the National Aeronautics and Space Administration, National Institutes of Health, and Michigan State University AgBioResearch.

<sup>2</sup> This manuscript resulted from the first author's presentation in the 59th Systematics Symposium "China," which was held at the Missouri Botanical Garden in St. Louis, Missouri, U.S.A., on 12–13 October 2012.

<sup>3</sup> Center for Systems Integration and Sustainability, Department of Fisheries and Wildlife, 1405 S. Harrison Road, Suite 115 Manly Miles Bldg., Michigan State University, East Lansing, Michigan 48823-5243, U.S.A. liuji@msu.edu, vina@msu.edu.

doi: 10.3417/2013040

many people inside and outside the protected areas still use them in a variety of ways, such as for harvesting timber, agriculture, mining, and tourism (Liu et al., 1999b). They are also exposed to the potentially negative effects of human-induced climate change (IPCC, 2007).

The world's biodiversity conservation challenge is well represented by China. Due to its enormous variation in climatic, topographic, and geologic features, China is one of the world's 17 mega-diverse countries (Mittermeier et al., 1997) and ranks third in the world (after Brazil and Colombia) in the number of known plant species (Liu et al., 2003b). China's flora is extremely rich and diversified, with around 33,000 vascular plant species in more than 3000 different genera and ca. 334 families of angiosperms, ca. 42 genera and 11 families of gymnosperms, 231 genera and 63 families of pteridophytes, and ca. 500 genera and 106 families of bryophytes (Lu, 2004; Wang, 2004; Wu et al., 2004a; Wu et al., 2004b). More than 10,000 higher plant species are endemic to China (Fu, 1992; Pitman & Jørgensen, 2002), with ca. 243 genera found only in China (Ying & Zhang, 1994). In addition, although not endemic, many plant genera have been recorded predominantly in China, including the 650 species of the genus *Rhododendron* L. (of the 1025 distributed worldwide) (Wu et al., 2003, 2004b). This high diversity is at high risk mainly due to deforestation (e.g., for timber and fuel wood), as well as the conversion of both forests and grasslands into croplands and urban areas. While China has been occupied by humans for millennia, the tremendous loss and fragmentation of natural ecosystems have been particularly drastic from the 1950s onward (Dinerstein & Wikramanayake, 1993; Liu, 2010). The huge destruction of China's ecosystems began to receive attention by government authorities due, in part, to the occurrence of natural disasters and the decrease in crop productivity associated with soil erosion and desertification. Therefore, starting in the 1970s, the government began to implement a series of afforestation and reforestation programs that have resulted in a progressive increase of forest cover over the last 40 years (Liu et al., 2013b). Yet nearly all of these new tree plantations, which are replacing previously natural forests, have been mono-specific and often consist of fast-growing exotic species (Viña et al., 2013) that greatly diminish the biodiversity value of the original forests (Xu & Wilkes, 2004). In addition, the government has established many protected areas at an impressive rate (Liu & Raven, 2010). By the end of 2011, a total of 2640 nature reserves were established covering ca. 14.9% of the land surface of

China (Ministry of Environmental Protection, People's Republic of China, 2011). Many endangered and charismatic species, such as the giant panda (*Ailuropoda melanoleuca* [David, 1869]), the South China tiger (*Panthera tigris* subsp. *amoyensis* [Hilzheimer, 1905]), and the Asian elephant (*Elaphas maximus* [Linnaeus, 1758]), have been used as surrogates for the establishment of these nature reserves. The giant panda is particularly interesting because it is considered an icon of biodiversity conservation not only in China but in the whole world (Mackinnon & De Wulf, 1994; Liu et al., 2001; Loucks et al., 2001).

Almost all ecosystems, including protected areas, constitute coupled human and natural systems (CHANS) (Liu et al., 2007a, 2007b) in which humans and natural components interact with each other directly or indirectly. Thus, understanding and conserving plants and wildlife require a new approach that treats humans, plants, and wildlife as parts of an integrated system. To illustrate the CHANS approach (Liu et al., 2007a, 2007b), in this article we focus on complex relationships among giant pandas, plants (the tree and bamboo species pandas depend on), and humans. We first outline a conceptual framework of CHANS and then highlight the major components and their interrelationships, using results mainly from our research group and collaborators.

#### CONCEPTUAL FRAMEWORK

The CHANS approach brings together theoretical foundations (e.g., social norms, social networks, ecological succession) and analytical techniques (e.g., remote sensing, geographic information systems, systems modeling) from diverse disciplines, including those from ecological and social sciences, to understand complex systems (Liu, 2001; Liu et al., 2007a; McConnell et al., 2011). The approach is, therefore, well suited for understanding pandas, the plants they depend on, and humans, which affect both pandas and plants. In this article, we conceptualize the integrated system as consisting of four main components: giant panda habitat, plants, local residents, and policies, all of which, in turn, are influenced by macroscale factors, including climate, natural disasters (e.g., earthquakes, landslides), and telecoupling processes that operate over distances (Liu et al., 2013a; Liu, 2014) (Fig. 1).

Each of the components in our framework both influences and is influenced by the other components. For instance, the abundance, distribution, and diversity of plants (i.e., tree and understory bamboo species) are influenced by people (e.g., local




Figure 1. Conceptual framework of coupled human and natural systems (CHANS) for synthesizing information on the relationships among pandas, the tree and bamboo species they depend on, and humans (including conservation policies). Arrows represent the direction of influences among different system components, with those going in both directions representing feedbacks. The dotted line represents the permeable boundary of CHANS.

residents) through different activities such as farming, cutting trees for timber or fuel wood, collection of bamboo shoots, and tree and bamboo plantation (State Forestry Administration, 2006). All these activities modify the habitat suitability of pandas, thus affecting their abundance, distribution, and behavior. While hunting panda individuals is strictly prohibited and the sanctions imposed are drastic enough to discourage this practice, some illegal hunting still occurs, although it is mostly unintended (e.g., pandas may be inadvertently captured by wild boar snares; Lü & Kemf, 2001). Nevertheless, humans have other direct effects on the abundance, distribution, and behavior of pandas, since human settlements pose obstacles to dispersal not only within nature reserves, but also among nature reserves and mountain regions (Xu et al., 2006). Therefore, humans and their activities are at odds with the survival of pandas and of the tree and bamboo species they depend on (Liu et al., 2004).

In turn, the changes in the abundance and distribution of plants and pandas provide feedback to humans through changes in the accessibility to timber and non-timber forest products (e.g., bamboo shoots) and motivate the development and imple-

mentation of conservation policies that act both directly (e.g., preventing timber extraction and fuel wood collection; spurring tree planting) and indirectly (e.g., incentives to use alternatives to fuel wood, such as electricity). These policies constitute feedback mechanisms through which changes in the abundance and distribution of plants and pandas may be reduced. In short, policies influence human attitudes and activities, which in turn change panda habitat. Changes in panda habitat may prompt the government to develop and implement new policies. These interrelationships result in numerous feedback loops.

Finally, the relations and feedbacks among the four main components in the framework (Fig. 1) are influenced by macroscale factors. These include changes in climate (IPCC, 2007; Tuanmu et al., 2013), the impact of natural disasters such as earthquakes (Viña et al., 2011; Zhang et al., 2011), and telecoupling processes (Liu & Yang, 2013) such as labor migration and tourism, which in many cases have opposing effects.

#### GIANT PANDAS

The historical distribution of the species at one time covered an area of ca. 2.2 million km<sup>2</sup>




Figure 2. Spatial distribution of the historical (gray) and current (green) geographic ranges of the giant panda (Mackinnon & De Wulf, 1994; Reid & Gong, 1999; Hu & Wei, 2004; State Forestry Administration, 2006; Viña et al., 2010). Also shown are the six mountain regions where pandas currently survive, as well as the 63 nature reserves (including the flagship Wolong Nature Reserve) established for the conservation of the pandas and their habitat.

throughout 19 provinces of China, northern Vietnam, and northern Myanmar (Schaller et al., 1985; Pan et al., 2001) (Fig. 2). However, by the early 1800s this distribution was reduced to only five provinces in southwestern China comprising ca. 262,000 km<sup>2</sup>, which was further reduced to ca. 124,000 km<sup>2</sup> by the early 1900s (Zhu & Long, 1983; Reid & Gong, 1999). The current distribution comprises only ca. 21,000 km<sup>2</sup> within six mountain regions (i.e., Qionglai, Minshan, Qionglai, Greater Xiangling, Lesser Xiangling, and Liangshan) in three provinces (Gansu, Shaanxi, and Sichuan) of southwestern China (Mackinnon & De Wulf, 1994; Reid & Gong, 1999; Hu & Wei, 2004; State Forestry Administration, 2006; Viña et al., 2010) (Fig. 2).

In terms of the total number of pandas, no estimates before the 1970s (when range-wide censuses started to be performed) are available, although expert opinion suggests that the population could have been ca. 3000 individuals during the 1950s (Hu, 2001). The first census in the 1970s estimated

that ca. 2400 individuals survived in the wild, and the second census in the 1980s estimated a decline of more than 50%, to ca. 1100 (Hu, 2001). The third census (performed between 2001 and 2003) estimated the total number of giant panda individuals at around 1600 (State Forestry Administration, 2006). The increase in the number of pandas between the second and the third censuses has been attributed mostly to larger regions surveyed and better techniques used to assess the number of individuals living in a region during the latter census, rather than to an actual increase in the panda population, although such an increase cannot be completely rejected (State Forestry Administration, 2006).

Pandas are extreme dietary specialists; 99% of their diet is composed of understory bamboo species (Schaller et al., 1985). Adaptations to a strict bamboo diet include an enlarged wrist bone to allow for gripping bamboo stems and a large skull and jawbone for mastication of tough plant material (Schaller et al., 1985). However, the panda's short digestive tract

(characteristic of a carnivore) together with a lack of gut microflora suitable for breaking down the cellulose contained in their diet allows the pandas to digest less than 20% of their food intake (Schaller et al., 1985). To respond to these digestive constraints, the pandas have developed behavioral adaptations such as maintaining low energy expenditures (e.g., pandas prefer areas with gentle slopes for ease of movement) and selecting different bamboo species and plant parts with different nutritional characteristics that change along the year (Schaller et al., 1985). Because understory bamboo species in the forests of southwestern China are located at particular elevation ranges, the selection of different bamboo species along the year makes pandas elevation migrants (Linderman et al., 2004). Since the understory bamboo species require forest cover for shade, and pandas also depend on forest cover for shelter (Liu et al., 1999b, 2001), there is a tight relationship between pandas, understory bamboo, and tree species.

#### PLANT SPECIES IN THE PANDA GEOGRAPHIC RANGE

The current geographic range of the giant panda is characterized by high mountains and deep valleys with elevations ranging from ca. 70 m to more than 6000 m. The drastic elevation gradient combined with complex terrain, geology, and soils is responsible for a high biological diversity, including more than 6000 species of plants in more than 1000 genera (IUCN, 2006). The region also features more than 100 species of mammals in 25 families, and ca. 400 species of birds in 45 families (Reid & Hu, 1991; Taylor & Qin, 1993b; IUCN, 2006). In fact, the region comprises one of the world's top 25 biodiversity hotspots (Myers et al., 2000; Mittermeier et al., 2004).

The vegetation is dominated by evergreen and deciduous broadleaf forests at intermediate and lower elevations and subalpine coniferous forests at higher elevations. Forests occupy around one third of the current panda range, with coniferous, broadleaf deciduous, and mixed coniferous/broadleaf deciduous forests accounting for ca. 48%, 32%, and 20% of the forest cover, respectively (Fig. 3). Tree species diversity in these forests is influenced by elevation and shows conspicuous declines in the number of tree species with an increase in elevation, particularly above 2500 m (Viña et al., 2012).

A total of 115 tree species were found in 104 field plots randomly distributed within an elevation range of 1000–3000 m in broadleaf deciduous, coniferous, and mixed forests across the Qinling Mountains region (Fig. 2) (Viña et al., 2012). Dominant tree

species (i.e., occurring in > 20% of the field plots) in this mountain region were *Quercus aliena* Blume, *Betula albosinensis* Burkill, *Prunus scopulorum* Koehne, *Toxicodendron vernicifluum* (Stokes) F. A. Barkley, and *Pinus armandii* Franch. (Viña et al., 2012). In a similar study in the Wolong Nature Reserve (the first and one of the largest giant panda nature reserves, established in 1975 and encompassing ca. 200,000 ha in Sichuan Province) located in the Qionglai Mountains region (Liu et al., 1999a) (Fig. 2), a total of 81 tree species were found in 62 field plots distributed along four transects ranging in elevation from 1500 to 3200 m (unpubl. data). Dominant tree species (i.e., occurring in > 20% of the field plots) in this region were *Acer laxiflorum* Pax, *Salix dolia* C. K. Schneid., *Prunus pilosiuscula* (C. K. Schneid.) Koehne, *B. albosinensis*, *Abies faxoniana* Rehder & E. H. Wilson, *Corylus ferox* Wall. var. *tibetica* (Batalin) Franch., and *Sorbaria arborea* C. K. Schneid.

Bamboo species are a particularly conspicuous feature in the understory of the forests of the panda geographic range. Although pandas can feed on more than 30 understory bamboo species found across their geographic range, they prefer only ca. 14 spatially dominant ones (Table 1). In the Qinling Mountains region the understory canopy is dominated by three bamboo species: *Fargesia qinlingensis* T. P. Yi & J. X. Shao, *Bashania fargesii* (E. G. Camus) Keng f. & T. P. Yi [= *Arundinaria fargesii* E. G. Camus], and *F. dracocephala* T. P. Yi (Viña et al., 2012), while in the Wolong Nature Reserve (Qionglai Mountains region), the dominant bamboo species are *F. nitida* (Mitford) Keng f. ex T. P. Yi, *F. robusta* T. P. Yi, and *B. faberi* (Rendle) T. P. Yi [= *A. faberi* Rendle] (Linderman et al., 2005b; Viña et al., 2008; Tuanmu et al., 2010). Because there is usually only one dominant understory bamboo species at a given location (Johnson et al., 1988; Reid et al., 1989; Linderman et al., 2005b, 2006; Viña et al., 2008; Tuanmu et al., 2010; Viña et al., 2012), changes in their abundance and spatial distribution (e.g., through bamboo harvesting; bamboo die-offs as a result of mass flowering occurring at time intervals from 16- to 90-year cycles depending on the species; cf. Table 1) could cause food shortages for the pandas (Schaller, 1987; Reid et al., 1989).

The spatial distribution of understory bamboo species is influenced by many factors, including panda utilization and interactions with other plant species. Regarding the former, Pleistocene fossil records (Zhu & Li, 1980) indicate that the giant panda and the bamboo species composing their diet had plenty of time to develop co-evolutionary traits




Figure 3. Distribution of coniferous, broadleaf deciduous, and mixed coniferous/deciduous forests in the mountain regions (in 2007) comprising the geographic range of the giant panda. Coniferous forests occur mostly at higher elevations, while mixed coniferous/deciduous forests and broadleaf deciduous forests occur more so at intermediate and lower elevations, respectively. Some coniferous forests also occur at intermediate elevations but are usually planted assemblages.

that ensured their survival, e.g., by reducing potentially negative effects of overgrazing. Nevertheless, under the current drastically reduced areas of suitable habitat, the degree to which bamboo overutilization potentially results in a decline in both quality of bamboo stands and viability of the panda populations needs to be assessed. Studies conducted in Wolong Nature Reserve have found that *Fargesia robusta* had a low yearly recruitment rate, reflecting poor shoot production and survival rates (Taylor & Qin, 1993a). This was in part due to the intensive utilization of this species by pandas, particularly between May and June when adult individuals were found to consume as much as 35 kg of bamboo per day (Schaller et al., 1985). Since panda habitat areas located at lower elevations are becoming increasingly fragmented and degraded over time due to human

activities (Liu et al., 2001; Xu et al., 2006; Viña et al., 2007), there is potential for localized panda populations to over-utilize stands of *F. robusta* in isolated areas (Hull et al., 2010). However, much more work on this regard needs to be conducted. Regarding the interactions between bamboo and other plant species, the occurrence of overstory tree species influences the occurrence of understory bamboo (Taylor et al., 2004, 2006). Multivariate statistical analyses can be used to identify overstory tree species associated with understory bamboo species. Using the Jaccard similarity index (Jaccard, 1908) combined with hierarchical cluster analyses (i.e., average linkage method), we determined the tree species most likely associated with the six dominant bamboo species in the two field study sites (i.e., three bamboo species in the Qinling Mountains region and

Table 1. Dominant understory bamboo species occurring in the different mountain regions across the entire giant panda range. Mountain regions are indicated in the second column as (1) Qinling; (2) Minshan; (3) Qionglai; (4) Xiangling (both the Greater and Lesser Xiangling); and (5) Liangshan. These bamboo species also constitute the preferred food consumed by pandas. Also shown in the third column are their reported reproductive cycles (i.e., periods between mass flowering events). Names noted in bracketed equivalencies reflect a differing taxonomic usage in the *Flora of China* (Wu et al., 2006).

Species	Mountain region	Reproductive cycle (yrs.)	References
<i>Bashania fargesii</i> (E. G. Camus) Keng f. & T. P. Yi [= <i>Arundinaria fargesii</i> E. G. Camus]	1, 2, 3	23	Schaller et al., 1989; Wang, 1989; Tian, 1990; Pan & Lu, 1993; Zhou & Pan, 1997; State Forestry Administration, 2006; Viña et al., 2012
<i>B. fabri</i> (Rendle) T. P. Yi [= <i>A. fabri</i> Rendle]	2, 3, 4, 5	45	Campbell & Qin, 1983; Campbell, 1984; Johnson et al., 1988; Reid et al., 1989; Wang, 1989; Tian, 1990; Reid & Hu, 1991; Reid et al., 1991; Taylor & Qin, 1993a; State Forestry Administration, 2006
<i>Chimonobambusa pachystachys</i> Hsueh & T. P. Yi	2, 3, 4	16	Campbell & Qin, 1983; Wang, 1989; State Forestry Administration, 2006
<i>Fargesia scabrata</i> T. P. Yi	2, 3	50	Campbell, 1984; Schaller et al., 1983; Wang, 1989; State Forestry Administration, 2006
<i>F. denudata</i> T. P. Yi	2	40	Schaller et al., 1989; Wang, 1989; State Forestry Administration, 2006
<i>F. dracocephala</i> T. P. Yi	1, 2	n/a	State Forestry Administration, 2006; Li et al., 2007; Viña et al., 2012
<i>F. fenax</i> (Keng) T. P. Yi	4, 5	40	Wang, 1989; State Forestry Administration, 2006
<i>F. nitida</i> (Mitford) Keng f. ex T. P. Yi	1, 2, 3	90	Wang, 1989; Tian, 1990; State Forestry Administration, 2006
<i>F. qinlingensis</i> T. P. Yi & J. X. Shao	1	50	Tian, 1987; State Forestry Administration, 2006; Wang et al., 2007; Viña et al., 2012
<i>F. robusta</i> T. P. Yi	1, 2, 3	37	Campbell & Qin, 1983; Campbell, 1984; Johnson et al., 1988; Reid et al., 1989; Wang, 1989; Tian, 1990; Reid & Hu, 1991; Reid et al., 1991; Pan & Lu, 1993; Zhou & Pan, 1997; State Forestry Administration, 2006
<i>F. rufa</i> T. P. Yi	2	15	Wang, 1989; State Forestry Administration, 2006
<i>Qiongzhuca opienensis</i> (J. R. Xue & T. P. Yi) D. Z. Li & J. R. Xue [= <i>Chimonobambusa opienensis</i> (J. R. Xue & T. P. Yi) T. H. Wen & Ohrnberger]	4, 5	20	Wang, 1989; State Forestry Administration, 2006
<i>Yushania chungii</i> (Keng) Z. P. Wang & G. H. Ye [= <i>Y. brevipaniculata</i> (Hand.-Mazz.) T. P. Yi]	1, 2, 3	20	Campbell & Qin, 1983; Campbell, 1984; Wang, 1989; State Forestry Administration, 2006
<i>Y. confusa</i> (McClure) Z. P. Wang & G. H. Ye	4, 5	22	Wang, 1989; State Forestry Administration, 2006



three in the Wolong Nature Reserve in the Qionglai Mountains region) (Fig. 4). In both of these regions the species *Betula albosinensis*, a dominant tree species in the giant panda geographic range, was associated with at least three dominant understory bamboo species (Fig. 4). Other species of the genus *Betula* L. (e.g., *B. utilis* D. Don, *B. platyphylla* Sukaczew, *B. luminifera* H. J. P. Winkl.) were also associated with some of the dominant bamboo species (Fig. 4). The occurrence of these species associations can be explained by the interactions between tree species life histories and bamboo life cycles which influence both tree and bamboo regeneration and, thus, contribute to structuring the species composition in the forests of the panda geographic range (Taylor et al., 2004).

#### PEOPLE

The pattern of decline of both the distribution and number of pandas has been largely associated with the loss of tree and understory bamboo species, which, in turn, have been attributed to climate fluctuations over many centuries (Schaller et al., 1985; Reid & Gong, 1999; Pan et al., 2001). However, the most recent loss has been attributed to an ever increasing growth of human population and activities (e.g., agricultural expansion, timber harvesting, road construction, livestock grazing, tourism, bamboo harvesting, mining) (Schaller et al., 1985; Reid & Gong, 1999; Pan et al., 2001), together with a drastic increase in the number of households. For example, from 1975 to 2000, the number of local residents in Wolong Nature Reserve increased by 72.4% (from 2560 to 4413) while the number of households jumped by 129.9% (from 421 to 968) (An et al., 2011). The increase in the number of households may have even more drastic negative effects than population growth alone (Liu et al., 2003a), because the proliferation of smaller households (e.g., as a result of divorces and kids moving out of parental homes [An et al., 2003]) translates into more households and less efficient use of natural resources on a per-capita basis (Yu & Liu, 2007).


Human activities vary across space and over time. For example, while cultivation has occurred in the panda geographic range for centuries, it mostly occurred at low elevations which were more accessible to humans (Pan et al., 2001; Loucks et al., 2003). However, in recent decades, cultivation has extended to higher elevations, thus, induced forest clearings where core panda habitat was previously located (Schaller et al., 1985; Wang, 2008). In addition, the extensive timber harvesting and fuel wood collection observed in the panda region

between the 1960s and the 1990s were identified to be the most important threats to pandas (Liu et al., 2001; State Forestry Administration, 2006), particularly because clear-cut areas are characterized by two extremes that prevent the occurrence of panda individuals: complete lack of bamboo or extremely dense patches of non-nutritious bamboo without an overstory of trees (Schaller et al., 1985; Bearer et al., 2008).

In addition to changes in the distribution and composition of tree and bamboo species, timber harvesting and fuel wood collection have many other effects that generate feedbacks which ultimately affect local residents. For example, inside Wolong Nature Reserve, timber harvesting occurred along a spatially defined deforestation front in which forests were exploited near households during the 1970s (He et al., 2009). As forests near households depleted, the local residents were forced to cut trees farther away (He et al., 2009). The spatial expansion of timber-extraction activities was also driven by improved accessibility due to the development of road networks. For instance, a project by the United Nations' World Food Programme in the early 1980s designed to mitigate human impacts on panda habitat in the Reserve resulted in a road expansion. This road expansion facilitated the access to a forested area rarely visited during the 1970s, which was subsequently deforested (Wolong Administration Bureau, 2004; He et al., 2009).

Timber harvesting and fuel wood collection not only affect current panda habitat, but also generate legacy effects on future panda habitat (An et al., 2005, 2006; Linderman et al., 2005a) through their conspicuous effects on plant species. For instance, in Wolong Nature Reserve, wood constitutes the main material for traditional house construction. In addition, a significant amount of fuel wood is consumed for cooking, heating, and stewing pig fodder, since pigs are the main animal protein source for local residents (An et al., 2001; Liu et al., 2004). Deciduous tree species such as *Betula albosinensis* and *B. utilis* are preferred for fuel wood, while species such as *Abies faxoniana* are preferred for construction. As mentioned above, these deciduous and evergreen tree species are dominant overstory canopy components in the forests that are suitable for the pandas, but currently they are rarely found in areas near local households, due to excessive harvesting through selective logging. As these species promote the regeneration of understory bamboo suitable for the giant panda (Taylor et al., 2004, 2006), the absence of these species in some areas reduces the likelihood of suitable panda habitat regeneration.

## Qinling Mountains Region



## Wolong Nature Reserve (Qionglai Mountains Region)




Figure 4. Cluster analysis showing tree and bamboo species associations of six dominant bamboo species (underlined), three of which are located in the Qinling Mountains region and three in the Wolong Nature Reserve located within the Qionglai Mountains region (see Fig. 2 for location). Numbers in the figure correspond to: (1) *Betula platyphylla* Sukaczew.; (2) *Quercus spinosa* David ex Franch.; (3) *Ulmus parvifolia* Jacq.; (4) *Pyrus betulifolia* Bunge; (5) *Cornus controversa* Hemsl.; (6) *Quercus liaotungensis* Koidz.; (7) *Betula luminifera* H. J. P. Winkl.; (8) *Fargesia qinlingensis* T. P. Yi & J. X. Shao; (9) *Juglans cathayensis* Dode; (10) *Toxicodendron vernicifluum* (Stokes) F. A. Barkley; (11) *Fargesia dracocephala* T. P. Yi; (12) *Pinus*

Current emerging threats from human activities on pandas and their habitat include the extensive investment in infrastructure (e.g., dams, highways, airports, roads, railroads, tourism facilities) under the West China Development Program. Between 2001 and 2006 alone, this program invested more than U.S. \$125 billion for development projects with the main goal of attracting businesses and migrants to western China (Lu, 2009). This construction boom may exacerbate the degradation and fragmentation of giant panda habitat not only in the present but also in the years to come.

An additional threat comes in the form of livestock grazing (Hull et al., 2011). The third panda census reported that livestock grazing is the second most commonly encountered human-driven disturbance, after deforestation (State Forestry Administration, 2006). In the Minshan Mountain region (see its location in Fig. 1), disturbances due to livestock grazing were found in 19% of over 1600 sample plots (Wang, 2008). Despite the recorded prevalence of livestock grazing across the panda geographic range, research on its direct and indirect effects on pandas and their habitat is limited to a handful of case studies (Ran, 2003, 2004; Ran et al., 2003; Kang et al., 2011). These studies mostly reinforced the findings about the prevalence of livestock disturbance in panda habitat areas, hinting that there is some degree of overlap in the habitat selection of pandas and livestock. Therefore, many questions still remain unanswered, particularly regarding whether livestock could threaten the long-term sustainability of understory bamboo, which prior to the increase in livestock grazing was not believed to be threatened by any animal competing with the pandas (Schaller et al., 1985).

#### POLICIES

The impacts of human activities on pandas have prompted the Chinese government to develop a series of policies, including the establishment of panda nature reserves (protected areas specifically designed to conserve the panda). A total of 63 nature reserves (by 2007) have been designated to conserve the habitat for the panda. While these panda reserves

contain about 40% of the current panda habitat (Viña et al., 2010), they tend to be isolated (Viña et al., 2007, 2010), which means that there is still a significant amount of panda habitat outside the nature reserve system. However, nature reserves have not been immune to human threats, as the panda habitat contained within their boundaries also experienced a steady decrease (Liu et al., 2001; Viña et al., 2007). This has occurred despite the fact that these reserves have imposed restrictions on resource use (e.g., fuel wood) (He et al., 2009) and are required to establish zoning designations to spatially contain human activities (Hull et al., 2011). In many cases these designations are ineffectual because the boundaries of zoning designations are displaced in response to development pressures (e.g., construction of tourism facilities) (Jim & Xu, 2004; Liu & Li, 2008; Hull et al., 2011). In addition, panda reserves often lack sufficient funding for their operations (Liu et al., 2003b) and, thus, do not properly enforce conservation activities (Lü & Kemf, 2001).

In addition, since the late 1990s the Chinese government has been implementing two of the largest ecological conservation programs in the world: (1) The Natural Forest Conservation Program (NFCP), which bans logging in natural forests in order to prevent illegal harvesting (Yang et al., 2013b), and (2) the Grain-to-Green Program (GTGP; also referred to as the Sloping Land Conversion Program), which encourages farmers to return steep cropland to forest or grassland (Uchida et al., 2005; Liu et al., 2008). Both of these policies have global implications, as they fulfill part of China's commitment to international biodiversity conservation treaties (Liu et al., 2008). These policies seem to have been producing overall positive effects on forests, including those in panda habitat regions (Li et al., 2013), by preventing further deforestation and promoting forest recovery (Viña et al., 2007, 2011; Liu et al., 2008, 2013). The implementation of the NFCP has started to also exhibit a significant positive effect on panda habitat, particularly when local people are actively involved in forest monitoring activities (Tuanmu, 2012). This is because timber harvesting has been curbed since

---

*armandii* Franch.; (13) *Quercus aliena* Blume; (14) *Betula albosinensis* Burkill; (15) *Prunus* L. sp.; (16) *Acer davidii* Franch.; (17) *Betula albosinensis* var. *septantrionalis* C. K. Schneid.; (18) *Abies fargesii* Franch.; (19) *Bashania fargesii* (E. G. Camus) Keng f. & T. P. Yi [= *Arundinaria fargesii* E. G. Camus]; (20) *Larix chinensis* Beissn.; (21) *Euonymus phellomanus* Loes.; (22) *Abies faxoniana* Rehder & E. H. Wilson; (23) *Fargesia nitida* (Mitford) Keng f. ex T. P. Yi; (24) *Betula utilis* D. Don; (25) *Prunus dielsiana* (C. K. Schneid.) Koehne; (26) *Quercus aquifolioides* Rehder & E. H. Wilson; (27) *Salix dissa* C. K. Schneid.; (28) *Salix phanera* C. K. Schneid.; (29) *Tilia intonsa* E. H. Wilson; (30) *Acer laxiflorum* Pax; (31) *Corylus ferox* Wall. var. *thibetica* (Batalin) Franch.; (32) *Fargesia robusta* T. P. Yi; (33) *Rhododendron pachytrichum* Franch.; (34) *Tsuga chinensis* (Franch.) E. Pritz.; (35) *Cotoneaster moupinensis* Franch.; (36) *Sorbaria arborea* C. K. Schneid.; (37) *Salix dolia* C. K. Schneid.; (38) *Syringa* L. sp.; (39) *Bashania fabri* (Rendle) T. P. Yi [= *Arundinaria fabri* Rendle]; (40) *Prunus pilosiuscula* (C. K. Schneid.) Koehne.

the late 1990s due, in part, to the national logging ban implemented through the NFCP (State Forestry Administration, 2006; Viña et al., 2007, 2011; Li et al., 2013). This ban has brought not only a reduction in the rate of loss of natural forests, but also forest regeneration in previously clear-cut areas (Viña et al., 2011; Li et al., 2013). Such forest regeneration brings some hope for panda conservation, since pandas have been observed to use secondary forests that are 30+ years old if adequate understory bamboo is present (Viña et al., 2007; Bearer et al., 2008).

Nationwide, the GTGP has converted 8.8 million ha of cropland into forest or grassland (Liu et al., 2013b). In the Qionglai Mountains region, this program provides seedlings for ca. 48 tree species, many of which are exotic (Viña et al., 2013) and do not constitute suitable tree species that promote bamboo regeneration, nor giant panda habitat recovery. Among the species planted under the GTGP, *Cunninghamia lanceolata* (Lamb.) Hook., *Magnolia officinalis* Rehder & E. H. Wilson, *Ligustrum lucidum* W. T. Aiton, *Cryptomeria japonica* (Thunb. ex L. f.) D. Don, *Eucommia ulmoides* Oliv., and *Alnus cremastogyne* Burkill are the species used more often (Viña et al., 2013). The Wolong Nature Reserve has, in addition to the GTGP (Chen et al., 2009a, 2009b, 2010, 2012b), a local Grain-to-Green/Bamboo Program (GTGB) in which farmers are compensated for actively planting the species *Fargesia robusta* in previous cropland areas (Yang et al., 2013a, 2013b). The planted bamboo is harvested particularly to provide fodder for the captive pandas in the breeding center of the Reserve. As these monospecific bamboo plantations generally lack overstory trees, they could not be considered suitable habitat areas for the pandas, either now or in the foreseeable future. Thus, promoting the re-growth of tree species that characterize suitable panda habitat should be one of the priorities in the panda conservation toolbox.

#### MACROSCALE FACTORS

Plants and pandas are not only affected by local factors, but also by large-scale factors such as migration, tourism, natural disasters, and climate change. Since China has been the fastest-growing economy in the world over the past three decades (Liu & Diamond, 2008), there has been an increase in job opportunities in cities; thus, more rural people are engaging in labor migration. This has the effect of reducing deforestation (e.g., through a reduction in fuel wood demands) while enhancing the natural regeneration of forests, particularly in giant panda regions (Chen et al., 2012a). In addition, many nature

reserves in China are exhibiting a rapid increase in the influx of tourists from around the world (Han, 2000; Liu et al., 2003a). The increase in tourism is supported by rapid development of tourism facilities (e.g., hotels, restaurants), together with expansions in the road network, all of which influence the abundance and distribution of plants and pandas. For example, the number of tourists visiting Wolong Nature Reserve increased dramatically, from ca. 20,000 in 1995 to ca. 100,000 in 2000 (Lindberg et al., 2003) and to more than 200,000 in 2006 (He et al., 2008). During this time period, the number of households participating in tourism activities also increased, yet households with more livelihood assets (e.g., income, education, social capital) were more likely to participate than less affluent households (Liu et al., 2012).

Another macroscale factor is represented by natural disasters. As in most mountain regions around the world, landslides are the most common natural disaster in the panda geographic range (Brabb & Harrod, 1989; Lee, 2005; Pradhan et al., 2006). More than 70% of catastrophic landslides are related to land use change (Reid et al., 2006; Huang, 2007; Rindfuss et al., 2008) and are triggered by severe rainfall events during the summer months (Li, 1989), and by a high seismic activity, with four earthquakes larger than 7.0 on the Richter scale occurring in the region during the last hundred years (Wang et al., 2008). The last such earthquake was the 7.9  $M_w$  (8.0 Richter scale) 12 May 2008 earthquake, the epicenter of which was located in Wenchuan County, Sichuan Province. Earthquake-induced landslides not only affect many forest areas (Viña et al., 2011) and giant panda habitat (Wang et al., 2008; Xu et al., 2009) (Fig. 5), but also induce significant losses to tree and shrub species richness (Zhang et al., 2011). Earthquake-induced landslides are so severe in many areas that they offset the gains in forest cover obtained through the implementation of conservation policies such as the NFCP (Viña et al., 2011).

A final macroscale factor potentially affecting plants and pandas is human-induced climate change. While deforestation and forest degradation are threatening the survival of about half of all bamboo species worldwide (Bystrakova & Kapos, 2006), climate change may present an additional significant threat. Many bamboo species are vulnerable to climate change because their unusual extended sexual reproduction intervals (from 10 to 120 years) (Janzen, 1976) and limited seed dispersal ability (Taylor et al., 1991) render them less capable of adjusting their distributions to the rapidly changing climate projected to occur within this century (IPCC,




Figure 5. Spatial distribution of panda habitat areas (modified from Liu et al., 2001; Viña et al., 2007) in Wenchuan County damaged by the 12 May 2008, Wenchuan Earthquake. Wolong and Caopo nature reserves are located entirely within Wenchuan County.

2007). In addition, bamboo species have limited vegetative dispersal ability (e.g., ca. 0.2–0.35 m/year for *Bashania fargesii* and *Fargesia robusta*) (Tian, 1989; Taylor & Qin, 1993a). Yet, despite this vulnerability of bamboo species to climate change, few studies have evaluated the potential effects of climate change on bamboo distribution and how these will cascade to pandas. A recent ensemble of bamboo distribution projections associated with multiple climate change projections and bamboo dispersal scenarios indicates that a substantial reduction in the distributional ranges of the three dominant bamboo species in the Qinling Mountains of China may occur by the end of the 21st century, with potentially drastic effects in the distribution of suitable giant panda habitat (Tuanmu et al., 2013). For instance, two contrasting climate change projections derived from two general circulation models (one developed

by the Commonwealth Scientific and Industrial Research Organisation, Australia [CSIRO-Mk2] and the other from a coupled model developed by the Center for Climate System Research and the National Institute for Environmental Studies, Japan [CCSR/NIES]) under the conservative B2 greenhouse gas emission scenario (which emphasizes local solutions to economic, social, and environmental sustainability) (IPCC, 2007), project reductions and distributional shifts of climatically suitable areas (CSA) of the three bamboo species by the end of the century (i.e., 2070–2099). Under the CSIRO-Mk2, there is a conspicuous projected shift of the CSA of the three species toward higher latitudes as compared with current conditions (Tuanmu et al., 2013) (Fig. 6). Under the CCSR/NIES, the CSA of *F. qinlingensis* is projected to disappear, as this species currently occurs at higher elevations, while the CSA of *B. fargesii* and *F.*




Figure 6. Current and projected (2070–2099) distributions of climatically suitable areas for the three dominant bamboo species in the Qinling Mountains region of China (after Tuanmu et al., 2013): *Fargesia dracocephala* (red), *F. qinlingensis* (blue), and *Bashania fargesii* (green). Their co-occurrences are indicated by the overlapping colors of the three species circles. The climate projections were derived from two global circulation models (one developed by the Commonwealth Scientific and Industrial Research Organisation, Australia [CSIRO-Mk2], and the other a coupled model developed by the Center for Climate System Research and the National Institute for Environmental Studies, Japan [CCSR/NIES]) under the B2 scenario of the IPCC Special Report on Emission Scenarios (IPCC-SRES). The black line corresponds to the political boundary of the Qinling Mountains region in Shaanxi Province, China. For its location within the panda geographic range, see Figure 2.

*dracocephala* are projected to experience a substantial reduction (Tuanmu et al., 2013) (Fig. 6). However, irrespective of the global circulation model used, the projected location of the CSA is distant from current bamboo distribution ranges, which may hinder potential range shifts of bamboo species (as well as pandas) in response to the projected change in the global climate (Tuanmu et al., 2013). This is particularly exacerbated by the long history of human activities surrounding the Qinling Mountains, which are not projected to substantially reduce in the foreseeable future.

## CONCLUSIONS

The current pace of biodiversity loss worldwide is intensifying the need for conservation actions that address not only species of conservation interest, but also their interactions with multiple other species, as well as with humans. Using the CHANS framework, this article has illustrated the complex relationships among pandas, plants (tree and bamboo species), and

humans. Our synthesis indicates that many human factors, including socioeconomic and demographic, exhibit reciprocal interactions with pandas and the plant species that comprise their habitat. The framework is particularly useful for uncovering key patterns and processes behind plant–animal interactions modified by human activities. Understanding of such patterns and processes is crucial for developing and evaluating policies that better reconcile the needs of an ever-growing human population with biodiversity conservation.

## Literature Cited

- An, L., J. G. Liu, Z. Y. Ouyang, M. Linderman, S. Q. Zhou & H. M. Zhang. 2001. Simulating demographic and socioeconomic processes on household level and implications for giant panda habitats. *Ecol. Modelling* 140: 31–49.
- An, L., A. Mertig & J. Liu. 2003. Adolescents' leaving parental home: Psychosocial correlates and implications for conservation. *Populat. & Environm.* 24: 415–444.
- An, L., M. Linderman, A. Shortridge, J. Qi & J. Liu. 2005. Exploring complexity in a human–environment system:

- An agent-based spatial model for multidisciplinary and multiscale integration. *Ann. Assoc. Amer. Geogr.* 95: 54–79.
- An, L., G. He, Z. Liang & J. Liu. 2006. Impacts of demographic and socioeconomic factors on spatio-temporal dynamics of panda habitats. *Biodivers. & Conservation* 15: 2343–2363.
- An, L., M. Linderman, G. He, Z. Ouyang & J. Liu. 2011. Long-term ecological effects of demographic and socioeconomic factors in Wolong Nature Reserve (China). Pp. 179–195 in R. P. Cincotta & L. J. Gorenflo (editors), *Human Population: Its Influences on Biological Diversity*. Springer-Verlag, Berlin.
- Baillie, J. E. M., C. Hilton-Taylor & S. N. Stuart (editors). 2004. 2004 IUCN Red List of Threatened Species: A Global Species Assessment. IUCN Publications Services Unit, Cambridge, United Kingdom.
- Bearer, S., M. Linderman, J. Huang, L. An, G. He & J. Liu. 2008. Effects of fuel wood collection and timber harvesting on giant panda habitat use. *Biol. Conservation* 141: 385–393.
- Brabb, E. E. & B. L. Harrod (editors). 1989. *Landslides: Extent and Economic Significance*. A. A. Balkema, Rotterdam.
- Bystrakova, N. & V. Kapos. 2006. Bamboo diversity: The need for a Red List review. *Biodiversity* 6: 12–16.
- Campbell, J. J. N. 1984. Giant panda conservation and bamboo forest destruction. Pp. 599–615 in T. N. Veziroglu (editor), *The Biosphere: Problems and Solutions*. Elsevier, Amsterdam.
- Campbell, J. J. N. & Z. Qin. 1983. Interactions of giant panda, bamboo and people. *J. Amer. Bamboo Soc.* 4: 1–35.
- Caro, T. M. 2002. Factors affecting the small mammal community inside and outside Katavi National Park, Tanzania. *Biotropica* 34: 310–318.
- Ceballos, G., A. Garcia & P. R. Ehrlich. 2010. The sixth extinction crisis: Loss of animal populations and species. *J. Cosmol.* 8: 1821–1831.
- Chen, X., F. Lupi, G. He & J. Liu. 2009a. Linking social norms to efficient conservation investment in payments for ecosystem services. *Proc. Natl. Acad. Sci. U.S.A.* 106: 11,812–11,817.
- Chen, X., F. Lupi, G. He, Z. Ouyang & J. Liu. 2009b. Factors affecting land reconversion plans following a payment for ecosystem service program. *Biol. Conservation* 142: 1740–1747.
- Chen, X., F. Lupi, A. Viña, G. He & B. Liu. 2010. Using cost-effective targeting to enhance the efficiency of conservation investments in payments for ecosystem services. *Conservation Biol.* 24: 1469–1478.
- Chen, X., K. A. Frank, T. Dietz & J. Liu. 2012a. Weak ties, labor migration, and environmental impacts: Toward a sociology of sustainability. *Organ. & Environm.* 25: 3–24.
- Chen, X., F. Lupi, L. An, R. Sheely, A. Viña & J. Liu. 2012b. Agent-based modeling of the effects of social norms on enrollment in payments for ecosystem services. *Ecol. Modelling* 229: 16–24.
- DeFries, R., A. Hansen, B. L. Turner, R. Reid & J. G. Liu. 2007. Land use change around protected areas: Management to balance human needs and ecological function. *Ecol. Applic.* 17: 1031–1038.
- Díaz, S., J. Fargione, F. S. Chapin III & D. Tilman. 2006. Biodiversity loss threatens human well-being. *PLoS Biol.* 4: e277.
- Dinerstein, E. & E. D. Wikramanayake. 1993. Beyond “hotspots”: How to prioritize investments to conserve biodiversity in the Indo-Pacific region. *Conservation Biol.* 7: 53–65.
- Fu, L. K. 1992. *China Plant Red Data Book: Rare and Endangered Plants*. Science Press, Beijing and New York.
- Gilliam, F. S. 2007. The ecological significance of the herbaceous layer in temperate forest ecosystems. *Bioscience* 57: 845–858.
- Han, N. (editor). 2000. *Analyses and Suggestions for Management Policies of China’s Protected Areas*. Scientific and Technical Documents Publishing House, Beijing.
- He, G. M., X. D. Chen, W. Liu, S. Bearer, S. Q. Zhou, L. Y. Q. Cheng, H. M. Zhang, Z. Y. Ouyang & J. G. Liu. 2008. Distribution of economic benefits from ecotourism: A case study of Wolong Nature Reserve for giant pandas in China. *Environm. Managem.* 42: 1017–1025.
- He, G. M., X. D. Chen, S. Bearer, M. Colunga, A. Mertig, L. An, S. Q. Zhou, M. Linderman, Z. Y. Ouyang, S. Gage, S. X. Li & J. G. Liu. 2009. Spatial and temporal patterns of fuel wood collection in Wolong Nature Reserve: Implications for panda conservation. *Landscape & Urb. Plan.* 92: 1–9.
- Hilton-Taylor, C. 2000. 2000 IUCN Red List of Threatened Species. IUCN/SSC. Gland, Switzerland, and Cambridge, United Kingdom.
- Hooper, D. U., F. S. Chapin III, J. J. Ewell, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naehm, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer & D. A. Wardle. 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecol. Monogr.* 75: 3–35.
- Hu, J. 2001. *Research on the Giant Panda*. Shanghai Science and Education Publishing House, Shanghai.
- Hu, J. & F. Wei. 2004. Comparative ecology of giant pandas in five mountain ranges of their distribution in China. Pp. 137–147 in D. Lindburg & K. Baragona (editors), *Giant Pandas: Biology and Conservation*. University of California Press, Berkeley.
- Huang, R. 2007. Large-scale landslides and their sliding mechanisms in China since the 20th century. *Chin. J. Rock Mech. & Engin.* 26: 433–454.
- Hull, V., A. Shortridge, B. Liu, S. Bearer, S. Zhou, H. Zhang, Z. Ouyang & J. Liu. 2010. The impact of giant panda foraging on bamboo dynamics in an isolated environment. *Pl. Ecol.* 212: 43–54.
- Hull, V., W. H. Xu, W. Liu, S. Q. Zhou, A. Viña, J. D. Zhang, M. N. Tuanmu, J. Y. Huang, M. Linderman, X. D. Chen, Y. Huang, Z. Y. Ouyang, H. M. Zhang & J. G. Liu. 2011. Evaluating the efficacy of zoning designations for protected area management. *Biol. Conservation* 144: 3028–3037.
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate Change 2007: Impacts, Adaptation and Vulnerability*. Cambridge University Press, New York.
- IUCN. 2001. *IUCN Red List Categories and Criteria, Version 3.1*. Prepared by the IUCN, Gland, Switzerland, and Cambridge, United Kingdom. <<http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria>>, accessed 9 May 2014.
- IUCN. 2006. *Sichuan World Heritage Evaluation Report # 20*. Prepared by the IUCN, Gland, Switzerland, and Cambridge, United Kingdom.

- IUCN, UNEP-WCMC. 2012. The World Database on Protected Areas (WDPA). United Nations Environment Programme (UNEP). World Conservation Monitoring Centre (WCMC), Cambridge, United Kingdom.
- Jaccard, P. 1908. Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaud. Sci. Nat.* 44: 223–270.
- Janzen, D. H. 1976. Why bamboos wait so long to flower. *Annual Rev. Ecol. Syst.* 7: 347–391.
- Jim, C. Y. & S. S. W. Xu. 2004. Recent protected-area designation in China: An evaluation of administrative and statutory procedures. *Geogr. J. (Peking)* 170: 39–50.
- Johnson, K. G., G. B. Schaller & J. Hu. 1988. Responses of giant pandas to a bamboo die-off. *Natl. Geogr. Res.* 4: 161–171.
- Kang, D., L. Zhao & G. Song. 2011. Competition relationship between giant panda and livestock in Wanglang National Nature Reserve, Sichuan. *J. Northeast Forestry Univ.* 39: 74–76.
- Lee, S. 2005. Application of logistic regression model and its validation for landslide susceptibility mapping using GIS and remote sensing data journals. *Int. J. Remote Sensing* 26: 1477–1491.
- Lepczyk, C. A., C. H. Flather, V. C. Radeloff, A. M. Pidgeon, R. B. Hammer & J. Liu. 2008. Human impacts on regional avian diversity and abundance. *Conservation Biol.* 22: 405–416.
- Li, D. Z., Z. Guo & C. Stapleton. 2007. *Fargesia dracocephala*. P. 93 in Z. Y. Wu, P. H. Raven & D. Y. Hong (editors), *Flora of China*, Vol. 22. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Li, T. 1989. Landslides: Extent and economic significance in China. Pp. 271–286 in E. E. Brabb & B. L. Harrod (editors), *Landslides: Extent and Economic Significance*. A. A. Balkema, Rotterdam.
- Li, Y., A. Viña, W. Yang, X. Chen, J. Zhang, Z. Ouyang, Z. Liang & J. Liu. 2013. Effects of conservation policies on forest cover change in giant panda habitat regions. *China. Land Use Policy* 33: 42–53.
- Lindberg, K., C. Tisdell & D. Xue. 2003. Ecotourism in China's Nature Reserves. Pp. 103–128 in A. A. Lew (editor), *Tourism in China*. Haworth Hospitality Press, New York.
- Linderman, M., J. Liu, J. Qi, L. An, Z. Ouyang, J. Yang & Y. Tan. 2004. Using artificial neural networks to map the spatial distribution of understory bamboo from remote sensing data. *Int. J. Remote Sensing* 25: 1685–1700.
- Linderman, M., L. An, S. Bearer, G. He, Z. Ouyang & J. Liu. 2005a. Modeling the spatio-temporal dynamics and interactions of households, landscape, and giant panda habitat. *Ecol. Modelling* 183: 47–65.
- Linderman, M., S. Bearer, L. An, Y. C. Tan, Z. Y. Ouyang & H. G. Liu. 2005b. The effects of understory bamboo on broad-scale estimates of giant panda habitat. *Biol. Conservation* 121: 383–390.
- Linderman, M. A., L. An, S. Bearer, G. M. He, Z. Ouyang & J. G. Liu. 2006. Interactive effects of natural and human disturbances on vegetation dynamics across landscapes. *Ecol. Applic.* 16: 452–463.
- Liu, J. 2001. Integrating ecology with human demography, behavior, and socioeconomics: Needs and approaches. *Ecol. Modelling* 140: 1–8.
- Liu, J. 2010. China's road to sustainability. *Science* 328: 50.
- Liu, J. 2014. Forest sustainability in China and implications for a telecoupled world. *Asia Pacific Policy Studies* 1: 230–250.
- Liu, J. & J. Diamond. 2008. Science and government: Revolutionizing China's environmental protection. *Science* 319: 37–38.
- Liu, J. & P. H. Raven. 2010. China's environmental challenges and implications for the world. *Crit. Rev. Environm. Sci. Technol.* 40: 823–851.
- Liu, J. & W. Yang. 2013. Integrated assessments of payments for ecosystem services programs. *Proc. Natl. Acad. Sci. U.S.A.* 110: 16,297–16,296.
- Liu, J., Z. Ouyang, Y. Tan, J. Yang & S. Zhou. 1999a. Changes in human population structure and implications for biodiversity conservation. *Populat. & Environm.* 21: 45–58.
- Liu, J., Z. Ouyang, W. Taylor, R. Groop, Y. Tan & H. Zhang. 1999b. A framework for evaluating effects of human factors on wildlife habitats: The case on the giant pandas. *Conservation Biol.* 13: 1360–1370.
- Liu, J., M. Linderman, Z. Ouyang, L. An & H. Zhang. 2001. Ecological degradation in protected areas: The case of Wolong Nature Reserve for giant pandas. *Science* 292: 98–101.
- Liu, J., G. C. Daily, P. R. Ehrlich & G. W. Luck. 2003a. Effects of household dynamics on resource consumption and biodiversity. *Nature* 421: 530–533.
- Liu, J., Z. Ouyang, S. L. Pimm, P. H. Raven, X. Wang, H. Miao & N. Han. 2003b. Protecting China's biodiversity. *Science* 300: 1240–1241.
- Liu, J., Z. Ouyang, H. Zhang, M. Linderman, L. An, S. Bearer & G. He. 2004. A new paradigm for panda research and conservation: Integrating ecology with human demography, behavior, and socioeconomics. Pp. 217–225 in D. Lindburg & K. Baragona (editors), *Giant Pandas: Conservation Priorities for the 21st Century*. University of California Press, Berkeley.
- Liu, J., T. Dietz, S. R. Carpenter, M. Alberti, C. Folke, E. Moran, A. N. Pell, P. Deadman, T. Kratz & J. Lubchenco. 2007a. Complexity of coupled human and natural systems. *Science* 317: 1513–1516.
- Liu, J., T. Dietz, S. R. Carpenter, C. Folke, M. Alberti, C. L. Redman, S. H. Schneider, E. Ostrom, A. N. Pell, J. Lubchenco, W. W. Taylor, Z. Ouyang, P. Deadman, T. Kratz & W. Provencher. 2007b. Coupled human and natural systems. *Ambio* 36: 639–649.
- Liu, J., S. X. Li, Z. Y. Ouyang, C. Tam & X. D. Chen. 2008. Ecological and socioeconomic effects of China's policies for ecosystem services. *Proc. Natl. Acad. Sci. U.S.A.* 105: 9477–9482.
- Liu, J., V. Hull, M. Batistella, R. DeFries, T. Dietz, F. Fu, T. W. Hertel, R. C. Izaurralde, E. F. Lambin, S. X. Li, L. A. Martinelli, W. J. McConnell, E. F. Moran, R. Naylor, Z. Y. Ouyang, K. R. Polenske, A. Reenberg, G. D. Rocha, C. S. Simmons, P. H. Verburg, P. M. Vitousek, F. S. Zhang & C. Q. Zhu. 2013a. Framing sustainability in a telecoupled world. *Ecol. & Soc.* 18(2): 26.
- Liu, J., Z. Ouyang, W. Yang, W. Xu & S. Li. 2013b. Evaluation of ecosystem service policies from biophysical and social perspectives: The case of China. Pp. 372–384 in S. A. Levin (editor), *Encyclopedia of Biodiversity*, 2nd ed. Academic Press, Waltham, Massachusetts.
- Liu, W., C. A. Vogt, J. Luo, G. He, K. A. Frank & J. Liu. 2012. Drivers and socioeconomic impacts of tourism participation in protected areas. *PLoS ONE*. doi:10.1371/journal.pone.0035420. <<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0035420>>, accessed 6 May 2014.



- Liu, X. & J. Li. 2008. Scientific solutions for the functional zoning of nature reserves in China. *Ecol. Modelling* 215: 237–246.
- Loucks, C. J., Z. Lu, E. Dinerstein, H. Wang, D. M. Olson, C. Q. Zhu & D. J. Wang. 2001. Giant pandas in a changing landscape. *Science* 294: 1465–1465.
- Loucks, C. J., Z. Lu, E. Dinerstein, D. J. Wang, D. L. Fu & H. Wang. 2003. The giant pandas of the Qinling Mountains, China: A case study in designing conservation landscapes for elevational migrants. *Conservation Biol.* 17: 558–565.
- Lu, D. 2009. The economic consequence of labor mobility in China's regional development. *Asian Econ. Pap.* 8: 85–114.
- Lu, S. G. 2004. Flora of pteridophyta in China. Pp. 78–94 in Z. Y. Wu & X. Chen (editors), *Flora Republicae Popularis Sinicae*, Vol. 1. Science Press, Beijing.
- Lü, Z. & E. Kemf. 2001. Giant Pandas in the Wild: A WWF Species Status Report. WWF International, Gland, Switzerland.
- Mackinnon, J. & R. De Wulf. 1994. Designing protected areas for giant pandas in China. Pp. 127–142 in R. I. Miller (editor), *Mapping the Diversity of Nature*. Chapman & Hall, London.
- Margules, C. R. & R. L. Pressey. 2000. Systematic conservation planning. *Nature* 405: 243–253.
- McConnell, W. J., J. D. A. Millington, N. J. Reo, M. Alberti, H. Asbjornsen, L. A. Baker, N. Brozović, L. E. Drinkwater, S. A. Drzyzga, J. Fragoso, D. S. Holland, C. A. Jantz, T. A. Kohler, H. D. G. Maschner, M. Monticino, G. Podestá, R. G. Pontius Jr., C. L. Redman, D. Sailor, G. Urquhart & J. G. Liu. 2011. Research on coupled human and natural systems (CHANS): Approach, challenges and strategies. *Bull. Ecol. Soc. Amer.* 92: 218–228.
- Meir, E., S. Andelman & H. P. Possingham. 2004. Does conservation planning matter in a dynamic and uncertain world? *Ecol. Letters* 7: 615–622.
- Ministry of Environmental Protection, People's Republic of China. 2011. China National Biodiversity Conservation Strategy and Action Plan. China Environmental Science Press, Beijing.
- Mittermeier, R. A., P. R. Gil & C. G. Mittermeier. 1997. Megadiversity: Earth's Biologically Wealthiest Nations. Conservation International (CI), Washington, D.C.
- Mittermeier, R. A., P. R. Gil, M. Hoffmann, J. D. Pilgrim, T. B. Brooks, C. G. Mittermeier, J. L. Lamoreux & G. A. B. da Fonseca. 2004. Hotspots Revisited: Earth's Biologically Richest and Most Endangered Ecoregions. CEMEX, Mexico City.
- Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca & J. Kent. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Nilsson, M.-C. & D. A. Wardle. 2005. Understorey vegetation as a forest ecosystem driver: Evidence from the northern Swedish boreal forest. *Frontiers Ecol. & Environm.* 3: 421–428.
- Pan, W. & Z. Lu. 1993. The giant panda. Pp. 140–145 in I. Stirling (editor), *Bears: Majestic Creatures of the Wild*. Rodale Press, Emaus, Pennsylvania.
- Pan, W., Z. Lü, D. Wang & H. Wang. 2001. The Opportunity for the Giant Panda to Exist. Peking University Press, Beijing.
- Parks, S. A. & A. H. Harcourt. 2002. Reserve size, local human density, and mammalian extinctions in U.S. protected areas. *Conservation Biol.* 16: 800–808.
- Pereira, H. M., G. C. Daily & J. Roughgarden. 2004. A framework for assessing the relative vulnerability of species to land-use change. *Ecol. Applic.* 14: 730–742.
- Pimm, S. L. & P. H. Raven. 2000. Biodiversity: Extinction by numbers. *Nature* 403: 843–845.
- Pitman, N. C. A. & P. M. Jørgensen. 2002. Estimating the size of the world's threatened flora. *Science* 298: 989.
- Pradhan, B., R. P. Singh & M. F. Buchroithner. 2006. Estimation of stress and its use in evaluation of landslide prone regions using remote sensing data. *Nat. Hazards Oceanogr. Processes Satellite Data* 37: 698–709.
- Ran, J. 2003. Habitat selection by giant pandas and grazing livestock in the Xiaoxiangling Mountains of Sichuan Province. *J. Sun Yat-sen Univ., Social Science Edition* 23: 2253–2259.
- Ran, J. 2004. A survey of disturbance of giant panda habitat in the Xiaoxiangling Mountains of Sichuan Province. *Acta Theriologica Sin.* 24: 277–281.
- Ran, J., S. Liu, H. Wang, Z. Sun, Z. Zeng & S. Liu. 2003. Effect of grazing on giant pandas' habitat in Yele Nature Reserve. *Acta Theriologica Sin.* 23: 288–294.
- Reid, D. G. & J. Hu. 1991. Giant panda selection between *Bashania fangi* bamboo habitats in Wolong Reserve, Sichuan, China. *J. Appl. Ecol.* 28: 228–243.
- Reid, D. G. & J. Gong. 1999. Giant panda conservation action plan. Pp. 241–254 in C. Servheen, S. Herrero & B. Peyton (editors), *Bears: Status Survey and Conservation Action Plan*. IUCN, Gland, Switzerland, and Cambridge, United Kingdom.
- Reid, D. G., J. C. Hu, D. Sai, W. Wei & Y. Huang. 1989. Giant panda (*Ailuropoda melanoleuca*) behavior and carrying-capacity following a bamboo die-off. *Biol. Conservation* 49: 85–104.
- Reid, D. G., A. H. Taylor, J. C. Hu & Z. S. Qin. 1991. Environmental influences on bamboo (*Bashania fangi*) growth and implications for giant panda conservation. *J. Appl. Ecol.* 28: 855–868.
- Reid, R. S., T. P. Tomich, J. Xu, H. Geist, E. F. Lambin, R. S. De Fries, J. G. Liu, D. Alves, B. Agbola, A. Chhabra, A. Mather, T. Veldcamp, K. Kok, M. van Noordwijk, D. Thomas, C. Palm & P. Verburg. 2006. Linking land change science and policy: Current lessons and future integration (Chapter 7). Pp. 175–204 in E. F. Lambin & H. Geist (editors), *Land-Use and Land-Cover Change: Local Processes and Global Impacts*. (Global Change—The IGBP Series). Springer, Berlin.
- Rindfuss, R. R., B. Entwisle, S. J. Walsh, L. An, N. Badenoch, D. G. Brown, P. Deadman, T. P. Evans, J. Fox, J. Geoghegan, M. Gutmann, M. Kelly, M. Linderman, J. G. Liu, G. P. Malanson, C. F. Mena, J. P. Messina, E. F. Moran, D. C. Parker, W. Parton, P. Prasartkul, D. T. Robinson, Y. Sawangdee, L. K. Vanwey & P. H. Verburg. 2008. Land use change: Complexity and comparisons. *J. Land Use Sci.* 3: 1–10.
- Sala, O. E., F. S. Chapin, J. J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L. F. Huenneke, R. B. Jackson, A. Kinzig, R. Leemans, D. M. Lodge, H. A. Mooney, M. Oesterheld, N. L. Poff, M. T. Sykes, B. H. Walker, M. Walker & D. H. Wall. 2000. Global biodiversity scenarios for the year 2100. *Science* 287: 1770–1774.
- Schaller, G. B. 1987. Bamboo shortage not only cause of panda decline. *Nature* 327: 562.
- Schaller, G. B., J. Hu, W. Pan & J. Zhu. 1985. *The Giant Pandas of Wolong*. University of Chicago Press, Chicago.

- Schaller, G. B., Q. Teng, K. G. Johnson, X. Wang, H. Shen & J. Hu. 1989. The feeding ecology of giant pandas and Asiatic black bears in the Tangjiahe Reserve, China. Pp. 212–241 in J. L. Gittleman (editor), *Carnivore Behavior, Ecology and Evolution*. Cornell University Press, Ithaca, New York.
- State Forestry Administration, P. R. China. 2006. Report of the Third National Giant Panda Census. Science Publishing House, Beijing.
- Taylor, A. H. & Z. Qin. 1993a. Structure and dynamics of bamboos in the Wolong Natural Reserve, China. *Amer. J. Bot.* 80: 375–384.
- Taylor, A. H. & Z. Qin. 1993b. Bamboo regeneration after flowering in the Wolong Giant Panda Reserve, China. *Biol. Conservation* 63: 231–234.
- Taylor, A. H., D. G. Reid, Z. S. Qin & J. C. Hu. 1991. Spatial patterns and environmental associates of bamboo (*Bashania fangiiana* Yi) after mass-flowering in south-western China. *Bull. Torrey Bot. Club* 118: 247–254.
- Taylor, A. H., J. Y. Huang & S. Q. Zhou. 2004. Canopy tree development and undergrowth bamboo dynamics in old-growth *Abies-Betula* forests in southwestern China: A 12-year study. *Forest Ecol. Managem.* 200: 347–360.
- Taylor, A. H., S. W. Jang, L. J. Zhao, C. P. Liang, C. J. Miao & J. Y. Huang. 2006. Regeneration patterns and tree species coexistence in old-growth *Abies-Picea* forests in southwestern China. *Forest Ecol. Managem.* 223: 303–317.
- Tian, X. 1987. Ecological observation on flowering and regeneration of bamboo forests in Mt. Qinling. *Bamboo Res.* 2: 39–43.
- Tian, X. 1989. Shooting and growth of *Bashania fargesii*. *J. Bamboo Res.* 8: 400–407.
- Tian, X. 1990. Studies of the food base of giant panda in Qinling Mountains. *Acta Theriologica Sin.* 10: 88–96.
- Tuanmu, M.-N. 2012. Spatiotemporal Dynamics of Giant Panda Habitat: Implications for Panda Conservation Under a Changing Environment. Ph.D. Dissertation, Michigan State University, East Lansing.
- Tuanmu, M.-N., A. Viña, S. Bearer, W. Xu, Z. Ouyang, H. Zhang & J. Liu. 2010. Mapping understory vegetation using phenological characteristics derived from remotely sensed data. *Remote Sensing Environm.* 114: 1833–1844.
- Tuanmu, M.-N., A. Viña, G. J. Roloff, W. Liu, Z. Ouyang, H. Zhang & J. Liu. 2011. Temporal transferability of wildlife habitat models: Implications for habitat monitoring. *J. Biogeogr.* 38: 1510–1523.
- Tuanmu, M.-N., A. Viña, J. A. Winkler, Y. Li, W. Xu, Z. Ouyang & J. Liu. 2013. Climate change impacts on understory bamboo species and giant pandas in China's Qinling Mountains. *Nature Clim. Change* 3: 249–253.
- Uchida, E., J. T. Xu & S. Rozelle. 2005. Grain for green: Cost-effectiveness and sustainability of China's conservation set-aside program. *Land Econ.* 81: 247–264.
- Viña, A., S. Bearer, X. Chen, G. He, M. Linderman, L. An, H. Zhang, Z. Ouyang & J. Liu. 2007. Temporal changes in giant panda habitat connectivity across boundaries of Wolong Nature Reserve, China. *Ecol. Applic.* 17: 1019–1030.
- Viña, A., S. Bearer, H. Zhang, Z. Ouyang & J. Liu. 2008. Evaluating MODIS data for mapping wildlife habitat distribution. *Remote Sensing Environm.* 112: 2160–2169.
- Viña, A., M.-N. Tuanmu, W. Xu, Y. Li, Z. Ouyang, R. DeFries & J. Liu. 2010. Range-wide analysis of wildlife habitat: Implications for conservation. *Biol. Conservation* 143: 1960–1963.
- Viña, A., X. D. Chen, W. J. McConnell, W. Liu, W. H. Xu, Z. Y. Ouyang, H. M. Zhang & J. G. Liu. 2011. Effects of natural disasters on conservation policies: The case of the 2008 Wenchuan earthquake, China. *Ambio* 40: 274–284.
- Viña, A., M.-N. Tuanmu, W. Xu, Y. Li, J. Qi, Z. Ouyang & J. Liu. 2012. Relationship between floristic similarity and vegetated land surface phenology: Implications for the synoptic monitoring of species diversity at broad geographic regions. *Remote Sensing Environm.* 121: 488–496.
- Viña, A., X. Chen, W. Yang, W. Liu, Y. Li, Z. Ouyang & J. Liu. 2013. Improving the efficiency of conservation policies with the use of surrogates derived from remotely sensed and ancillary data. *Ecol. Indicators* 26: 103–111.
- Vitousek, P. M., H. A. Mooney, J. Lubchenco & J. M. Melillo. 1997. Human domination of Earth's ecosystems. *Science* 277: 494–499.
- Waltert, M., A. Mardiatuti & M. Muhlenberg. 2004. Effects of land use on bird species richness in Sulawesi, Indonesia. *Conservation Biol.* 18: 1339–1346.
- Wang, D., S. Li, S. Sun, H. Wang, A. Chen, S. Li, J. Li & Z. Lu. 2008. Turning earthquake disaster into long-term benefits for the panda. *Conservation Biol.* 22: 1356–1360.
- Wang, H. S. 2004. Flora of gymnosperm in China. Pp. 95–102 in Z. Y. Wu & X. Chen (editors), *Flora Republicae Popularis Sinicae*, Vol. 1. Science Press, Beijing.
- Wang, M. 1989. A Comprehensive Survey Report on China's Giant Panda and Its Habitat. Ministry of Forestry, P. R. China and WWF-World Wildlife Fund for Nature, Beijing.
- Wang, W., S. B. Franklin & J. R. Ouellette. 2007. Clonal regeneration of an arrow bamboo, *Fargesia qinlingensis*, following giant panda herbivory. *Pl. Ecol.* 192: 97–106.
- Wang, X. 2008. The Impact of Human Disturbance on Giant Panda Habitat in the Minshan Mountains. Chinese Academy of Sciences, Beijing.
- Wolong Administration Bureau. 2004. The Chronicle of Wolong Nature Reserve. Sichuan Science and Technology Press, Chengdu.
- Wu, C. L., S. Shi & Y. P. Zhang. 2004a. A case for conservation. *Nature* 428: 213–214.
- Wu, C. Y., P. H. Raven & D. Y. Hong (editors). 2006. *Flora of China*, Vol. 22. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Wu, Z. Y., A. M. Lu, Y. C. Tang, Z. D. Chen & D. Z. Li. 2003. The Families and Genera of Angiosperms in China: A Comprehensive Analysis. Science Press, Beijing.
- Wu, Z. Y., H. H. Peng, D. Z. Li, Z. K. Zhou & H. Sun. 2004b. Flora of angiosperm in China. Pp. 121–583 in Z. Y. Wu & X. Chen (editors), *Flora Republicae Popularis Sinicae*, Vol. 1. Science Press, Beijing.
- Xu, J. & A. Wilkes. 2004. Biodiversity impact analysis in northwest Yunnan, southwest China. *Biodivers. & Conservation* 13: 959–983.
- Xu, W., Z. Ouyang, A. Viña, H. Zheng, J. Liu & Y. Xiao. 2006. Designing a conservation plan for protecting the habitat for giant pandas in the Qionglai mountain range, China. *Diversity & Distrib.* 12: 610–619.
- Xu, W., X. Wang, Z. Ouyang, J. Zhang, Z. Li, Y. Xiao & H. Zheng. 2009. Conservation of giant panda habitat in

- 
- South Minshan, China, after the May 2008 earthquake. *Frontiers Ecol. & Environm.* 7: 353–358.
- Yang, W., W. Liu, A. Viña, J. Luo, G. He, Z. Ouyang, H. Zhang & J. Liu. 2013a. Performance and prospects of payments for ecosystem services programs: Evidence from China. *J. Environm. Managem.* 127: 86–95.
- Yang, W., W. Liu, A. Viña, M.-N. Tuanmu, G. He, T. Dietz & J. Liu. 2013b. Nonlinear effects of group size on collective action and resource outcomes. *Proc. Natl. Acad. Sci. U.S.A.* 110: 10,916–10,921.
- Ying, T. S. & Y. L. Zhang. 1994. *The Endemic Genera of Seed Plants in China*. Science Press, Beijing.
- Yu, E. & J. Liu. 2007. Environmental impacts of divorce. *Proc. Natl. Acad. Sci. U.S.A.* 104: 20,629–20,634.
- Zhang, J., V. Hull, W. Xu, J. Liu, Z. Ouyang, J. Y. Huang, X. Wang & R. Li. 2011. Impact of the 2008 Wenchuan earthquake on biodiversity and giant panda habitat in Wolong Nature Reserve, China. *Ecol. Res.* 26: 523–531.
- Zhou, Z. H. & W. S. Pan. 1997. Analysis of the viability of a giant panda population. *J. Appl. Ecol.* 34: 363–374.
- Zhu, J. & Y. Li. 1980. *The Giant Panda*. Science Press, Beijing.
- Zhu, J. & Z. Long. 1983. The vicissitudes of the giant panda. *Acta Zool. Sinica* 29: 93–104.