



Forest Carbon and Climate Program
Department of Forestry
MICHIGAN STATE UNIVERSITY



State and Tribal Capacity Building on Forest Carbon

Forest Carbon and Climate Change in New Jersey

This technical briefing summarizes topics such as forest densities and cover types, carbon storage, and climate considerations for the state of New Jersey.

This technical briefing was made possible by funding from Penn Soil Resource Conservation and Development Council under a cooperative agreement with the U.S. Department of Agriculture, Forest Service.



EASTERN REGION

Table of Contents


New Jersey Forest Overview	2
Temperature and Precipitation	3
Projected Future Trends in Temperature / Precipitation.....	4
Forest Density	5
Forest Cover Types and Carbon.....	6
Forest Carbon Pools.....	7
Forest Carbon Density.....	8
Species-Specific Considerations for Climate Adaptation.....	9
Habitat Suitability and Migration Models	10
Adaptability Ratings.....	11
Climate Change Atlas Summary for Pitch Pine.....	11
Citations:	12

New Jersey Forest Overview

New Jersey is situated along the east coast of the United States and lies within the US Forest Service's Eastern Region (USFS Region 9). Bordering states include Pennsylvania to the west, New York to the northeast, and Delaware to the southwest, with the Atlantic coast marking the majority of New Jersey's southern and eastern boundaries.

A map of percent tree canopy cover in New Jersey is shown in **Figure 1**. This state has variable forest coverage across its extent, with the highest levels of canopy cover occurring in the northernmost portion of the state coinciding with higher elevations of the Appalachian Mountain range. Another area of consistent canopy cover exists in the inland southern portion of the state coinciding with New Jersey's Pine Barrens. Areas of reduced coverage coincide with major transportation corridors and areas of higher population densities near urban centers.

Figure 1. Percent tree canopy cover in New Jersey.



Temperature and Precipitation

Two major factors affecting forest carbon and productivity are temperature and precipitation. **Figure 2** shows normal mean temperatures throughout New Jersey between 1991 and 2020. Over this 30-year period, mean annual temperatures varied by about 9 °F across this state. Temperature trends largely follow latitudinal gradients, with warmer mean temperatures occurring in the southern portions of the state and giving way to cooler temperatures to the north. The warmest mean annual temperature is around 57 °F and occurs in the southwest and southernmost tip of New Jersey, while the coolest mean annual temperature is around 48 °F in the northwest portion of the state and coincides with higher elevations.

Figure 2. Normal mean temperature (°F) from 1991–2020 in New Jersey.




Figure 3. Normal mean precipitation (in.) from 1991-2020 in New Jersey.





Figure 3 shows normal mean precipitation throughout New Jersey between 1991 and 2020 and demonstrates the geographic variation in these trends. Over this 30-year period, mean annual precipitation levels varied by about 8 in. Areas that receive the lowest levels of precipitation (45-47 in.) occur along the state's southern boundary. Areas receiving the highest amounts of precipitation (52-53 in.) occur in the northwest mountainous portion of the state.

Projected Future Trends in Temperature / Precipitation

Figure 4. Model results for potential changes in temperature and precipitation trends in New Jersey through 2099 under a high emission scenario (RCP 8.5).

Potential Changes in Climate Variables



Projected future trends in temperature and precipitation for New Jersey between 2009 and 2099 are shown in **Figure 4**. Model results suggest average temperatures will continue to increase through the end of the century, a trend which is also projected for the coldest and warmest month averages, as well as throughout the growing season (May – Sep.). Over this 90-year period, average annual temperatures are expected to increase by an estimated 9.5 °F, with the most drastic increases expected to occur during the growing season (+10.2 °F).

Model results of future precipitation in New Jersey follow similar trends, with totals projected to steadily increase through 2099 (**Figure 4**). Over a 90-year period, annual precipitation is expected to increase by an estimated 9 in., which is a higher rate of change than projections for the growing season (+2.9 in.). This suggests that the most significant changes to precipitation in New Jersey may occur during the winter months (Oct. – Apr.).



Forest Density

Figure 5. Forest density as live tree density (No. ha⁻¹) in New Jersey.

Figure 6. Forest density as live tree basal area (m² ha⁻¹) in New Jersey.

Forest Density: Live tree number

Forest Density: Live tree basal area



Forest density¹ is both a structural characteristic of forests and a reflection of forest dynamics. It can be measured as the number of trees per unit area, or it can be measured in terms of live tree area per unit area, known as “basal area”. Live tree basal area represents the amount of ground covered by living trees in two-dimensional space. **Figure 5** shows average forest density in terms of live trees per hectare by ecosection² across the state of New Jersey, while **Figure 6** represents forest density by ecosection in terms of basal area (m² ha⁻¹).

By comparing these figures we can see that the large ecosection in the southeastern portion of New Jersey has the state’s highest forest density in terms of number of trees per hectare (**Figure 5**), but an average density in terms of basal area (**Figure 6**). This suggests that in this zone, there may be many total trees per unit area, but these trees tend to be relatively small. By contrast, an ecosection in the northern portion of the state that touches New Jersey’s western border has the state’s lowest forest density in terms of number of trees but has the highest forest density in terms of basal area, suggesting the prevalence of fewer, relatively large trees in this zone.

¹All forest inventory and carbon data were estimated using data from the Forest Inventory and Analysis (FIA) Program which can be accessed through the FIA DataMart (USDA Forest Service, 2024. *Forest inventory and analysis program*. Available at: <https://www.fia.fs.usda.gov/>) using the rFIA package (Stanke et al, 2020. rFIA: an R package for estimation of forest attributes with the US Forest Inventory and analysis database. *Environ Model Softw.* 127:104664. <https://doi.org/10.1016/j.envsoft.2020.104664>) in the R programming environment (R Core Team, 2020. *R: A language and environment for statistical computing*, Vienna, Austria: R Foundation for Statistical Computing.

²Ecosection definition can be found at Cleland et al, 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. *General Technical Report WO-76D*, Washington Office, USDA Forest Service. <https://doi.org/10.2737/WO-GTR-76D>

Forest Cover Types and Carbon

Figure 7. Total forest area (thousand ha) by forest type³ in New Jersey.





Figure 8. Total forest carbon (million tons) by forest type in New Jersey. Total forest carbon is the sum of carbon stored across all aboveground and belowground pools (Soil Organic carbon + Live Belowground carbon + Live Aboveground carbon + Litter carbon + Dead wood carbon).




New Jersey is dominated by 6 key forest cover types: Oak / hickory, Loblolly / shortleaf pine, Oak / pine, Oak / gum / cypress, Maple / beech / birch, and Elm / ash / cottonwood. **Figure 7** and **Figure 8** show state-level data of total forested area and total forest carbon, respectively, for each of these cover type groups. As these figures show, Oak / hickory is the dominant forest type of New Jersey, spanning an area close to 200,000 hectares and storing roughly 55 million tons of carbon statewide. With coverage levels ranging from ~30,000-180,000 hectares, other forest types in this state are less abundant, yet play an important role contributing to enhanced biodiversity and landscape heterogeneity. Comparing trends from **Figure 7** with those in **Figure 8** demonstrates how carbon storage levels vary by forest cover type. For example, Oak / pine forests cover slightly more land area than Oak / gum / cypress stands in New Jersey (**Figure 7**), yet when it comes to carbon, Oak / gum / cypress stands store slightly more carbon than their Oak / pine counterparts (**Figure 8**).

³Forest Types are a classification of forest land based upon and named for the tree species that forms the plurality of live-tree stocking. These forest types used in the briefing align with FIA's definition of Forest type group which are a combination of forest types that share closely associated species and site requirements. Longer definitions of both forest types and forest type groups are found in Appendix D of the Forest Inventory and Analysis Database: Database Description and User Guide for Phase 2 (version 9.1) which can be accessed here: https://research.fs.usda.gov/sites/default/files/2023-11/wo-fiadb_user_guide_p2_9-1_final.pdf

Forest Carbon Pools

Figure 9. Total forest carbon (million tons) by pool and forest type in New Jersey.




Forest carbon storage can be further assessed by examining how it's distributed across different ecosystem carbon pools. **Figure 9** shows the amount of carbon stored in different carbon pools of key forest cover types in New Jersey. These values show how different forest types allocate distinct proportions of forest carbon into soil organic matter, live belowground (BG) biomass, live aboveground (AG) biomass, litter, and dead wood pools. New Jersey forests generally allocate more ecosystem carbon to belowground pools (soil organic matter + live BG biomass) than aboveground pools (live AG biomass + litter + dead wood), yet the proportions in which they do so varies across forest cover types. For instance, Elm / ash / cottonwood forests allocate roughly 2x the amount of carbon to belowground pools than aboveground pools, whereas forest types like Oak / hickory and Maple / beech / birch distribute carbon more evenly between belowground and aboveground pools. Another noteworthy trait shown in **Figure 9** is the magnitude of carbon storage levels across different pools and cover types. Oak / hickory's dominating presence on this landscape means its statewide carbon pools are outsized compared to other groups. For example, leaf litter and dead wood pools of New Jersey's Oak / hickory forests on their own contain roughly the same levels of stored carbon as the total ecosystem carbon (sum of carbon stored across all pools) contained by the Elm / ash / cottonwood group.

Forest Carbon Density

Figure 10. Aboveground live forest carbon density (tC ha^{-1}) by forest type in New Jersey.

Average Forest Carbon Density by Ecoregion: Aboveground Live



Forest carbon density can be influenced by many ecosystem traits, such as tree density, stand age, species mix/ cover type, soil fertility, elevation, and a site's management and disturbance history. In **Figure 10**, the carbon density of aboveground living forest biomass is shown for 6 key cover types in New Jersey. Of these, Oak / pine stands hold the highest levels of aboveground live carbon per unit area, represented by the deep shade of green in a northwestern ecoregion. By contrast, Elm / ash / cottonwood stands have a much lower carbon density per unit area in this ecoregion. Across much of their extent, Loblolly / shortleaf pine and Elm / ash / cottonwood stands exhibit relatively even carbon densities, while cover types like Oak / pine and Maple / beech / birch show higher levels of variability across ecoregions. In these instances, variable carbon densities can be driven by the relative prevalence or absence of each forest type from a given ecoregion.

Species-Specific Considerations for Climate Adaptation

Climate change is expected impact the distribution of species into the future. Predictive modeling of potential future changes that incorporate species interactions, dispersal mechanisms, demography, physiology, and evolution is needed to assist in adaptive forest planning. The USDA Forest Service **Climate Change Tree Atlas, Version 4**, provides modeled potential suitable habitat for 125 species in the eastern US, with an additional 23 species. <https://www.fs.usda.gov/nrs/atlas/tree/>

Core Climate Change Atlas components:

- DISTRIB-II: Species habitat suitability model
- SHIFT: Migration model (when combined with DISTRIB-II, estimates colonization potential (HQCL) of future suitable habitats)
- Adaptability Ratings: Species adaptability ratings (species traits not included in DISTRIB-II and SHIFT models)

In addition to the modeled potential suitable habitat for individual tree species, the Climate Change Atlas includes Current and potential future habitat, capability and migration for individual tree species and potential changes in climate variables summarized by the following spatial extents:

Geographic Area	Description
National Forest Summaries	Results summarized for 55 national forests
National Park Summaries	Results summarized for 78 national parks
HUC6 Watershed	Results summarized by hydrologic unit codes level 3 (HUC 6) which are hierarchical classifications based on surface hydrologic features in which level 3 maps watershed basins (Seaber et al, 1987) https://pubs.usgs.gov/wsp/wsp2294/
Ecoregional Vulnerability Assessments (EVAS)	Results summarized by ecoregions used in the USDA Climate Hub Regional Vulnerability Assessments https://www.climatehubs.usda.gov/assessments
USDA Forest Service EcoMap 2007 Sections	Results summarized by ecological sections that delineate ecosystems with distinctive vegetation and other unique ecological characteristics (Cleland et al, 2007, McNab et al, 2007)
National Climate Assessment (NCA) 2015 Regional Summaries	Results summarized by National Climate Assessment Region which include the Midwest, Northeast, Northern Plains, Southeast, and Southern Plains
1 x 1° Grid Summaries	Results summarized by 1x1° latitude and longitude
State Summaries	Results summarized for 38 states
Urban areas	Results summarized for 185 urban areas across the eastern US

Additional background on this tool can be found at: <https://research.fs.usda.gov/centers/ccrc> along with short video tutorials on the Climate Change Atlas website.


Habitat Suitability and Migration Models

Model Reliability: **High**

Key Species Example: Modeled potential suitable habitat for Sugar Maple (*Acer saccharum*) through 2100


Current habitat quality and distribution (DISTRIB-II)

Potential migration (SHIFT) and colonization likelihood (CL)




Importance value is a measure of abundance that accounts for both tree basal area and number of stems, ranging from 0-100.

Colonization potential of future habitats under a high emission scenario (RCP 8.5)




Colonization is limited to range margins and infill (Blue) which is derived from habitat quality (DISTRIB-II) and migration model (SHIFT) utilizing the colonization likelihood model (CL). Orange shading represents current species' distributions where abundance is predicted to decrease due to loss of habitat suitability.

DISTRIB-II + SHIFT: Habitat quality and colonization likelihood (RCP 8.5)



Adaptability Ratings

Key Species Example: Pitch Pine (*Pinus rigida*)



V Hi Pos +3	High Pos +2	Low Pos +1	Minimal 0	Low Neg -1	High Neg -2	V Hi Neg -3
----------------	----------------	---------------	--------------	---------------	----------------	----------------

The Adaptability score, which assesses 21 variables to assign adaptability ratings to tree species in the eastern US, reflects a species' potential adaptability to climate change-driven stressors and disturbances at range wide scale. Adaptability ratings provide broad insights into factors that cannot be directly included in the Climate Change Tree Atlas species migration models. Two types of species traits are evaluated: 1) biological and 2) disturbance, each with their own set of factors to help characterize species' traits and responses to disturbance. Uncertainty is also included for each trait or factor assessed. When coupled with other modeled projections, adaptability ratings can support future planning under a changing climate.

The Adaptability variable is single score derived from the Modification Factors which encompass scores for the 12 disturbance and 9 biological factors. The Adaptability results can be considered relative to other tree species. For example, a species with a low Adaptability variable likely does not have life history characteristics to allow it to thrive under most conditions whereas a high Adaptability variable will likely do better under the climate change outputs from the DISTRIB-II and SHIFT Models.

Climate Change Atlas Summary for Pitch Pine

Pitch pine is a narrow distributed (1.9% of area), sparse, but of high importance were found. It was modeled to show a slight decrease in habitat (less so under RCP 8.5). It is moderate in abundance and adaptability with a fairly reliable model, yielding an overall capability to cope of fair for RCP 8.5 and poor for RCP 4.5. SHIFT migration is not significant.

Citations:

Habitat suitability models on trees:

Peters et al. (2020). Climate change tree atlas, Version 4. U.S. Forest Service, Northern Research Station and Northern Institute of Applied Climate Science, Delaware, OH. <https://www.nrs.fs.fed.us/atlas>;

Iverson, L.R, Peters, M.P., Prasad, A.M., & Matthews, S.N. (2019). Analysis of Climate Change Impacts on Tree Species of the Eastern US: Results of DISTRIB-II Modeling. *Forests*, 10(4), 302. doi: 10.3390/f10040302 <https://www.fs.usda.gov/treearch/pubs/57857>

Peters, M. P., Iverson, L. R., Prasad, A. M., & Matthews, S. N. (2019). Utilizing the density of inventory samples to define a hybrid lattice for species distribution models: DISTRIB-II for 135 eastern U.S. trees. *Ecology and Evolution*. doi: 10.1002/ece3.5445 <https://www.fs.usda.gov/treearch/pubs/58353>

Iverson, L. R., Prasad, A. M., Peters, M. P., & Matthews, S. N. (2019). Facilitating Adaptive Forest Management under Climate Change: A Spatially Specific Synthesis of 125 Species for Habitat Changes and Assisted Migration over the Eastern United States. *Forests*, 10(11), 989. doi: 10.3390/f10110989 <https://www.fs.usda.gov/treearch/pubs/59105>

Prasad, A. M., Iverson, L. R., Matthews, S. N., & Peters, M. P. (2016). A multistage decision support framework to guide tree species management under climate change via habitat suitability and colonization models, and a knowledge-based scoring system. *Landscape Ecology*, 31(9), 2187–2204. doi: 10.1007/s10980-016-0369-7 <https://www.fs.usda.gov/treearch/pubs/50748>

Prasad, A. M., Gardiner, J. D., Iverson, L. R., Matthews, S. N., & Peters, M. (2013). Exploring tree species colonization potentials using a spatially explicit simulation model: implications for four oaks under climate change. *Global Change Biology*, 19(7), 2196–2208. doi: 10.1111/gcb.12204 <https://www.fs.usda.gov/treearch/pubs/43705>

Iverson, L. R., A. M. Prasad, S. N. Matthews, and M. Peters. 2008. Estimating potential habitat for 134 eastern US tree species under six climate scenarios. *Forest Ecology and Management* 254:390-406. <http://www.treearch.fs.fed.us/pubs/13412>

Adaptability of tree species:

Iverson, L. R., S. N. Matthews, A. M. Prasad, M. P. Peters, et al. (2012). Development of risk matrices for evaluating climatic change responses of forested habitats. *Climatic Change* 114(2): 231-243. doi: 10.1007/s10584-012-0412-x. <https://www.fs.usda.gov/treearch/pubs/41221>

Matthews, S. N., L. R. Iverson, A. M. Prasad, M. P. Peters, and P. G. Rodewald. 2011. Modifying climate change habitat models using tree species-specific assessments of model uncertainty and life history factors. *Forest Ecology and Management* 262:1460-1472. <http://www.fs.usda.gov/treearch/pubs/38643>

Climate summary definitions:

McNab, W.H.; Cleland, D.T.; Freeouf, J.A.; Keys, Jr., J.E.; Nowacki, G.J.; Carpenter, C.A., comps. 2007. Description of ecological subregions: sections of the conterminous United States [CD-ROM]. Gen. Tech. Report WO-76B. Washington, DC: U.S. Department of Agriculture, Forest Service. 80 p. <https://research.fs.usda.gov/treearch/48669>

Cleland, D.T.; Freeouf, J.A.; Keys, J.E.; Nowacki, G.J.; Carpenter, C.A.; and McNab, W.H. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on CD-ROM] (A.M. Sloan, cartographer). Washington, DC: U.S. Department of Agriculture, Forest Service, presentation scale 1:3,500,000; colored. <https://research.fs.usda.gov/treearch/48672>

Seaber, Paul R., F. Paul Kapanos, and George L. Knapp (1987). Hydrologic Unit Maps. United States Geological Survey Water-Supply Paper 2294: i–iii, 1–63.