

# grapes.msu.edu

[Home](#)[About Us](#)[Search](#)[Newsletters](#)[Weather /  
Climate](#)[Viticulture](#)[Pest  
management](#)[Scouting guide](#)[Publications](#)[Calendar of  
events](#)[Industry links](#)[Contacts](#)

Download [Adobe Acrobat Reader](#) to view pdf files.

## Simazine injury

Tom Zabadal , MSU Horticulture

[Home](#) > [Scouting guide](#)> simazine

Injury from this herbicide appears as yellowing in the leaves between veins that remain green. In more advanced stages, portions or the entire leaf becomes brown. As with diuron, simazine injury tends to occur on light soils and sites where erosion has exposed roots near the soil surface.

### Additional information

- Search [MSU Extension News for Agriculture](#) site
- Search [MSU Fruit CAT Alert](#) newsletter for articles
- [MSU Diagnostic Services](#)



Simazine herbicide injury showing the characteristic yellowing between green veins. As the injury becomes more severe, leaves become lighter, eventually becoming brown and falling from the vine. Photo: T. Zabadal

[Site map](#)[Copyright/Linking](#)

Funding for this web site provided by [Project GREEN](#), [American Farmland Trust](#), [EPA Region 5's Strategic Agricultural Initiative](#) program, [The National Foundation for IPM Education](#), the [Center for Agricultural Partnerships](#) and the [MSU Integrated Pest Management Program](#) in collaboration with [MSU Extension](#) and the [Michigan Agricultural Experiment Station](#). Partially support from [NC-IPM Center](#).

05/26/11 Contact: [E. Haney](#)